

BRILL-NOETHER THEORY FOR CURVES ON GENERIC ABELIAN SURFACES

AREND BAYER AND CHUNYI LI

ABSTRACT. We completely describe the Brill-Noether theory for curves in the primitive linear system on generic abelian surfaces, in the following sense: given integers d and r , consider the variety $V_d^r(|H|)$ parametrizing curves C in the primitive linear system $|H|$ together with a torsion-free sheaf on C of degree d and $r + 1$ global sections. We give a necessary and sufficient condition for this variety to be non-empty, and show that it is either a disjoint union of Grassmannians, or irreducible. Moreover, we show that, when non-empty, it is of expected dimension.

This completes prior results by Knutsen, Lelli-Chiesa and Mongardi.

CONTENTS

| | |
|--|----|
| 1. Introduction | 1 |
| 2. Background: stability conditions, moduli spaces | 4 |
| 3. The Gieseker-wall | 8 |
| 4. Main proof | 11 |
| 5. Conclusion | 17 |
| 6. Generalisation | 20 |
| References | 21 |

1. INTRODUCTION

By Lazarsfeld's celebrated result [[Laz86](#)], a smooth curve in the primitive linear system of a generic K3 surface is Brill-Noether general (in the strongest possible sense, see [[Bay16](#), Theorem 1.1]). The corresponding question for abelian surfaces is much more subtle. In the present article, we completely determine when the Brill-Noether locus for the entire primitive linear system on generic abelian surfaces is non-empty, and show that, when non-empty, it is of the expected dimension. We show that the Brill-Noether locus is either irreducible or a disjoint union of Grassmannians. In particular, unlike for K3 surfaces our condition provides many examples with negative Brill-Noether numbers where *some* of the curves in the primitive linear system are not Brill-Noether general, in the sense of carrying torsion-free sheaves of prescribed degree and number of global sections. This completes previous work by Knutsen, Lelli-Chiesa and Mongardi [[KLCM15](#)].

Let X be an abelian surface whose Néron-Severi group $\mathrm{NS}(X)$ is generated by the class of an ample line bundle H . Let $g = \frac{H^2}{2} + 1$ be the arithmetic genus of curves in $|H|$. Given integers

$r \geq 1$ and $d \geq 1$, write $\chi = d + 1 - g$, and recall that the Brill-Noether number is given by $\rho(r, d, g) = g - (r + 1)(r + 1 - \chi)$. Let $V_d^r(|H|)$ denote the Brill-Noether locus parametrizing curves C in the linear system $|H|$ together with a pure sheaf L supported on C with $c_1(L) = H$, with $\chi(L) = \chi$, and $h^0(L) = r + 1$; this includes, of course, smooth curves in $|H|$ equipped with a complete linear system $g_d^r: C \rightarrow \mathbb{P}^r$ of degree d . Our main result is the following:

Theorem 1.1. *Assume $\chi \neq 0$. The Brill-Noether locus $V_d^r(|H|)$ is non-empty if and only if*

$$\rho + g - 2 \geq D|\chi| - D^2,$$

where D denotes the remainder of division of $r + 1$ by $|\chi|$. Moreover, when it is non-empty, it is generically smooth and of expected dimension $\rho + g - 2$.

When the above inequality is strict, then $V_d^r(|H|)$ is irreducible. Otherwise, it is a disjoint union of $\left(\frac{g-1}{\chi}\right)^2$ Grassmannians.

In [KLCM15, Theorems 1.4 and A.1], the authors showed that the last condition is necessary for non-emptiness, and that in this case the Brill-Noether locus has a component of expected dimension; they also showed that this condition is sufficient when $d \geq r(r + 1)$, and further implies the existence of *smooth* curves with g_d^r . In other words, in addition to their results we show that this necessary condition is also sufficient for $d < r(r + 1)$, and we determine when the Brill-Noether locus is irreducible.

In fact, a similar statement holds for certain moduli spaces of vector bundles. Let $\mathbf{v} \in H^*(X)$ be a class of the form $\mathbf{v} = (R, c_1(H), \chi)$ for some integers $R \in H^0(X, \mathbb{Z}) = \mathbb{Z}$ and $\chi \in H^4(X, \mathbb{Z}) = \mathbb{Z}$. Then the moduli space $M_H(\mathbf{v})$ of Gieseker-stable sheaves with Chern character v is smooth and irreducible of dimension $\mathbf{v}^2 + 2$. Let $M_H^{r+1}(\mathbf{v})$ denote the subset of sheaves E with $h^0(E) = r + 1$.

Theorem 1.2. *Assume that $\chi < 0$, and let r, g, D be as above. Then $M_H^{r+1}(\mathbf{v})$ is non-empty if and only if*

$$\mathbf{v}^2 - (r + 1)(r + 1 - \chi) \geq D(-\chi) - D^2.$$

In this case, it is irreducible and of expected dimension $\mathbf{v}^2 + 2 - (r + 1)(r + 1 - \chi)$.

1.1. Comparison with [KLCM15]. When $H^2 = 54$, in other word, $g = 28$, the following table lists the existence and emptiness result for curves in $V_d^r(|H|)$ for $20 \leq d \leq 26$ and $r \geq 1$.

| degree \ r(section) | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
|---------------------|----|-----|-----|--------|----------|--------|--------|
| 20 | BN | KLM | KLM | ϕ | ϕ | ϕ | ϕ |
| 21 | BN | BN | KLM | ϕ | ϕ | ϕ | ϕ |
| 22 | BN | BN | KLM | KLM | ϕ | ϕ | ϕ |
| 23 | BN | BN | KLM | KLM | ϕ | ϕ | ϕ |
| 24 | BN | BN | BN | KLM | Thm. 1.1 | ϕ | ϕ |
| 25 | BN | BN | BN | KLM | Δ | ϕ | ϕ |
| 26 | BN | BN | BN | KLM | Δ | ϕ | ϕ |

Table: (non-)emptiness of $V_d^r(|H|)$.

The labels in each box indicate the following situations:

ϕ : There is no line bundle of degree d with $r + 1$ section on a curve in $|H|$; this follows from [KLCM15, Theorem A.1] or our Theorem 1.1.

BN: The Brill-Noether number, $\rho(g, r, d)$, is non-negative, and thus every smooth curve carries a g_d^r by [Kem71, KL72],

KLM: We have $\rho(g, r, d) < 0$, but there exists a smooth curve with a g_d^r by [KLCM15, Theorem 1.4].

Δ : $V_d^r(|H|)$ is non-empty, see discussion below.

Thm. 1.1: The non-emptiness of $V_d^r(|H|)$ follows from Theorem 1.1.

We now give more explanation on the cells marked ‘ Δ ’. First of all, by our Theorem 1.1, $V_d^r(|H|)$ is non-empty. On the other hand, the non-emptiness of such $V_d^r(|H|)$ can be deduced from [KLCM15, Theorem 1.6 (i)] with the same arguments as that in [KLCM15, Example 5.15]. For example, when $r = 5$ and $d = 25$, one may let $k = 2$ and $\delta = 15$. It is direct to check that δ , k , and g (or p in the notation of [KLCM15]) satisfy the inequality in Theorem 1.6(i) in [KLCM15]. By the theorem, there is a curve C in $|H|$ with 15 nodes, and its normalization \tilde{C} is hyper-elliptic. In particular, \tilde{C} carries linear series with degree 10 and rank 5. Push-forward to C produces a torsion free sheaf with degree $10 + 15 = 25$ and rank 5. For the box that is marked by ‘Thm. 1.1’, the non-emptiness of $V_d^r(|H|)$ is due to Theorem 1.1 and is completely new. Interesting readers may check that Theorem 1.6 (i) in [KLCM15] does not provide suitable nodal curves that may carry linear series as desired.

Remark 1.3. Note that $V_d^r(|H|)$ in our setting is different from the space $|L|_d^r$ defined in [KLCM15]: for the latter, the support curve is required to be smooth.

1.2. Proof strategy. We proof Theorem 1.1 by wall-crossing. Let $\mathbf{v} = (0, H, \chi)$; then the moduli space $M_H(\mathbf{v})$ contains all pure torsion sheaves F with $c_1(F) = H$ and $\chi(F) = \chi$. It can be reinterpreted as the moduli space $M_\sigma(\mathbf{v})$ of Bridgeland-stable objects when σ is contained in the Gieseker-chamber for \mathbf{v} . We then show (for $\chi < 0$) that the first wall bounding the Gieseker-chamber destabilizes exactly those F with $h^0(F \otimes L_0) \neq 0$ for degree zero line bundle L_0 on X .

In particular, there is a natural short exact sequence

$$\mathcal{O}_X \otimes H^0(F) \hookrightarrow F \twoheadrightarrow F'$$

of semistable objects that is part of the Jordan-Hölder filtration of F . However, unlike in the case of K3 surfaces treated in [Bay16], we may have $h^0(F') \neq 0$; in fact, this may be necessary for the extension F to exist, as otherwise $\text{Ext}^1(F', \mathcal{O}_X)$ might be too small.

Thus, we have to proceed by induction. This induction is possible since all steps remain part of the finite Jordan-Hölder filtration of F . On the other hand, it is precisely this induction process that leads to the slightly arithmetic nature of our results (involving the division by remainder).

1.3. Generalisation. All our results hold slightly more generally for any polarized abelian surface (X, H) satisfying the following:

Assumption (*): H^2 divides $H \cdot D$ for all curve classes D on X .

To simplify the presentation, we first explain our entire argument in the case of Picard rank one; we then explain in Section 6 how to extend the arguments to this situation.

1.4. Related questions. In addition to [KLCM15], there have been a number of recent articles studying Brill-Noether loci of curves in M_g carrying a g_d^r for negative Brill-Noether numbers, in particular [Pfl13, Pfl17, JR17]. However, our results have no direct implication on this question, as we cannot prove that smooth curves appear in our locus $V_d^r(|H|)$.

1.5. Acknowledgments. It is a pleasure to thank Emanuel Macrì, Giovanni Mongardi, Soheyla Feyzbakhsh, and Sam Payne for a number of helpful discussions and comments. Both authors are supported by ERC starting grant no. 337039 “WallXBirGeom”.

2. BACKGROUND: STABILITY CONDITIONS, MODULI SPACES

2.1. Review: stability conditions on abelian surfaces. Let X be an abelian surface of Picard rank one; we denote by $D^b(X)$ the bounded derived category of coherent sheaves on X . In this section, we will review the description of a component of the space $\text{Stab}(X)$ of stability conditions on $D^b(X)$ given in [Bri08, Section 15].

Given an object $E \in D^b(X)$ we write $\text{ch}(E) \in H^*(X, \mathbb{Z})$ for its Chern character with value in cohomology. We write $H_{\text{alg}}^*(X, \mathbb{Z})$ for its algebraic part, i.e. the image of ch .

Let H be a line bundle as above; by abuse of notation, we will also write H for its class in $\text{NS}(X)$. Given $\beta \in \mathbb{R}$, we defined the β -twisted Chern character by

$$\text{ch}^\beta(E) := e^{-\beta H} \cdot \text{ch}(E) \in H_{\text{alg}}^*(X, \mathbb{R})$$

and the β -twisted slope of a coherent sheaf $E \in \text{Coh } X$ by

$$\mu_{H, \beta}(E) := \begin{cases} \frac{H \cdot \text{ch}_1^\beta(E)}{H^2 \text{ch}_0(E)} & \text{if } \text{ch}_0(E) > 0 \\ +\infty & \text{if } \text{ch}_0(E) = 0. \end{cases}$$

This leads to the usual notion of slope-stability, and the construction of the following torsion pair

$$\begin{aligned}\mathcal{T}^\beta &:= \langle E : E \text{ is } \mu_{H,\beta}\text{-semistable with } \mu_{H,\beta}(E) > 0 \rangle, \\ \mathcal{F}^\beta &:= \langle E : E \text{ is } \mu_{H,\beta}\text{-semistable with } \mu_{H,\beta}(E) \leq 0 \rangle,\end{aligned}$$

where $\langle \cdot \rangle$ denotes the extension-closure. This is a torsion pair in $\text{Coh } X$. Following [HRS96, Bri08], this lets us define a new heart of a bounded t-structure in $D^b(X)$ as follows:

$$\text{Coh}^\beta X := \langle \mathcal{F}^\beta[1], \mathcal{T}^\beta \rangle = \left\{ E : H^{-1}(E) \in \mathcal{F}^\beta, H^0(E) \in \mathcal{T}^\beta, H^i(E) = 0 \text{ for } i \neq 0, -1 \right\}.$$

For $\alpha > 0$ and $\beta \in \mathbb{R}$ as before, we define the central charge $Z_{\alpha,\beta} : K(X) \rightarrow \mathbb{C}$ by

$$(1) \quad Z_{\alpha,\beta}(E) := - \int_X e^{-i\alpha H} \cdot \text{ch}^\beta(E) = -\text{ch}_2^\beta(E) + i\alpha H \cdot \text{ch}_1^\beta(E) + \frac{\alpha^2}{2} H^2 \text{ch}_0(E).$$

Note that $Z_{\alpha,\beta}$ factors via the Chern character

$$(2) \quad \text{ch} : K(X) \rightarrow H_{\text{alg}}^*(X, \mathbb{Z}) \cong \mathbb{Z}^3.$$

We will first state Bridgeland's result constructing stability conditions on $D^b(X)$, and then explain its meaning.

Theorem 2.1 ([Bri08, Section 15]). *For $\alpha > 0, \beta \in \mathbb{R}$, the pair $\sigma_{\alpha,\beta} := (\text{Coh}^\beta X, Z_{\alpha,\beta})$ defines a stability condition on $D^b(X)$ satisfying the support property. Moreover, the map $\mathbb{R}_{>0} \times \mathbb{R} \rightarrow \text{Stab}(X)$ is continuous.*

We refer also to [MS16] or [Bay16] for more details and a sketch of the proof. Up to an action of the universal cover of $\text{GL}_2^+(\mathbb{R})$, the above theorem in fact describes an entire component of $\text{Stab}(X)$, but that fact will be irrelevant for us.

For our purposes, Theorem 2.1 makes two statements. First, consider the slope function

$$\nu_{\alpha,\beta} : \text{Coh}^\beta X \rightarrow \mathbb{R} \cup \{+\infty\}, \quad \nu_{\alpha,\beta}(E) := \begin{cases} \frac{-\Re Z_{\alpha,\beta}(E)}{\Im Z_{\alpha,\beta}(E)} & \text{if } \Im Z_{\alpha,\beta}(E) > 0 \\ +\infty & \text{if } \Im Z_{\alpha,\beta}(E) = 0. \end{cases}$$

This defines a notion of slope-stability in $\text{Coh}^\beta X$: every object admits a Harder-Narasimhan filtration into slope-semistable objects; every slope-semistable object admits a Jordan-Hölder filtration into slope-stable objects of the same slope.

Moreover, that these satisfy the support property just follows from the fact that on an abelian surface, the Chern character \mathbf{v} of any stable object satisfies $\mathbf{v}^2 \geq 0$; see Section 2.2 for a discussion of this fact.

The second statement, the continuity, implies that this family satisfies wall-crossing as α, β vary. To give an efficient description of the walls, we change our viewpoint slightly. Observe that up to the action of $\text{GL}_2(\mathbb{R})$ on $\mathbb{C} \cong \mathbb{R}^2$, a central charge $Z : K(X) \rightarrow \mathbb{C}$ that factors via the Chern character (2) is uniquely determined by its kernel $\text{Ker } Z \subset H_{\text{alg}}^*(X, \mathbb{R}) \cong \mathbb{R}^3$. The Mukai pairing:

$$\langle (v_0, v_1, v_2), (w_0, w_1, w_2) \rangle := v_1 w_1 - v_0 w_2 - v_2 w_0$$

equips $H_{\text{alg}}^*(X, \mathbb{R})$ with a quadratic form Q of signature $(2, 1)$. The definition of $Z_{\alpha, \beta}$ in equation (1) identifies the upper half plane $\mathbb{R}_{>0} \times \mathbb{R}$ with the projectivisation of the negative cone of Q , via the correspondence $(\alpha, \beta) \mapsto \text{Ker } Z_{\alpha, \beta}$.

Proposition 2.2 ([Bri08, Proposition 9.3]). *Let $\mathbf{v} \in H_{\text{alg}}^*(X, \mathbb{Z})$ be a primitive algebraic cohomology class. Then there exists a collection of two-dimensional linear subspaces $\mathcal{W}_v^i \subset H_{\text{alg}}^*(X, \mathbb{R}) \cong \mathbb{R}^3$ containing \mathbf{v} with the following properties:*

- *there exists a strictly $\sigma_{\alpha, \beta}$ -semistable object with Chern character \mathbf{v} if and only if $\text{Ker } Z_{\alpha, \beta}$ is contained in one of the subspaces \mathcal{W}_v^i ;*
- *the intersection of these subspaces with the negative cone of Q is locally finite; and*
- *as α, β vary within a chamber, i.e. without crossing a wall, (semi-)stability with respect to $\sigma_{\alpha, \beta}$ is unchanged for objects of Chern character \mathbf{v} .*

In the case where $\mathbf{v}^2 = 0$, the set of walls is empty.

See Figure 1 for a picture; we also refer to [Mac12, Theorem 3.1] for a proof of the fact that the walls are nested when viewed as semi-circles in the upper half plane.




FIGURE 1. Describing walls via $\text{Ker } Z_{\alpha, \beta} \subset H^*(X, \mathbb{R})$

The final claim, for the case $\mathbf{v}^2 = 0$, is a general fact for stability conditions satisfying the support property with respect to a given quadratic form, see e.g. [BMS16, Proposition A.8].

2.2. Moduli spaces and large-volume limit. Recall that on an abelian surface, there are no rigid objects (see [Bri08, Lemma 15.1] for a short proof). Therefore, if the moduli space $M_{\sigma_{\alpha,\beta}}(\mathbf{v})$ of $\sigma_{\alpha,\beta}$ -stable objects of Chern character $\mathbf{v} \in H^*(X, \mathbb{Z})$ is non-empty, then Hirzebruch-Riemann-Roch shows $\mathbf{v}^2 \geq 0$. It turns out that this necessary condition is also sufficient.

Theorem 2.3 ([Yos12, MYY11]). *Let $v \in H_{\text{alg}}^*(X, \mathbb{Z})$ be a primitive algebraic cohomology class. If α, β are generic, then the moduli space $M_{\sigma_{\alpha,\beta}}(\mathbf{v})$ of semistable objects of Chern character \mathbf{v} is non-empty if and only if $\mathbf{v}^2 \geq 0$. In this case it is a smooth holomorphic symplectic variety¹ and has dimension $\mathbf{v}^2 + 2$.*

Proof. In the case of $\mathbf{v}^2 > 0$, this is part of the statement of [Yos12, Theorem 1.13] and [MY11, Proposition 5.16]. (The statement “deformation-equivalent to ...” in [MY11, Proposition 5.16] in particular includes the non-emptiness.)

As for $\mathbf{v}^2 = 0$, in this case there are no walls as observed in Proposition 2.2. Therefore, we may assume that σ is near the large-volume limit, in which case it is well-known that the Gieseker-moduli space $M_{\sigma}(\mathbf{v}) = M_H(\mathbf{v})$ is an abelian surface derived equivalent to X . \square

Now fix such a primitive cohomology class \mathbf{v} of positive rank. If β remains fixed, and $\alpha \rightsquigarrow +\infty$, then the phase of $Z_{\alpha,\beta}(E)$ is asymptotically governed by the slope $\mu_H(E)$. The following result will refine that observation.

We write $M_H(\mathbf{v})$ for the moduli space of Gieseker-stable sheaves with respect to the polarization H . Given $E \in \text{D}^b(X)$, let $E^\vee = \mathbf{R}\mathcal{H}om(E, \mathcal{O}_X)$ be its derived dual, and write \mathbf{v}^\vee for the class dual to \mathbf{v} .

Proposition 2.4 ([Bri08, Proposition 14.2]). *If $\beta < \mu_H(\mathbf{v})$ and if α is sufficiently big, then $M_{\sigma_{\alpha,\beta}}(\mathbf{v}) = M_H(\mathbf{v})$: an object $E \in \text{Coh}^\beta(X)$ of Chern character \mathbf{v} is $\sigma_{\alpha,\beta}$ -stable if and only if it is a Gieseker-stable sheaf.*

If $\beta > \mu_H(\mathbf{v})$ and α is sufficiently big, then $M_{\sigma_{\alpha,\beta}}(-\mathbf{v}) = \mathbb{D}(M_H(\mathbf{v}^\vee))[1]$: an object $E \in \text{Coh}^\beta(X)$ of Chern character $-\mathbf{v}$ is $\sigma_{\alpha,\beta}$ -stable if and only if it is the shift $E = F^\vee[1]$ of the derived dual of a Gieseker-stable sheaf F of class \mathbf{v}^\vee .

In other words, there is a Gieseker-chamber in which Bridgeland stability and Gieseker stability coincide for objects of Chern character \mathbf{v} ; and similarly there is a dual Gieseker chamber in which they coincide up to taking derived duals.

We also make the following observation for non-primitive classes of square zero:

Proposition 2.5. *Let $\mathbf{v} = m\mathbf{v}_0$ with $m > 1$ and $\mathbf{v}_0^2 = 0$. Then every σ -semistable object of class \mathbf{v} is strictly semistable, and all its Jordan-Hölder factors are of class \mathbf{v}_0 .*

In particular, such an object has a stable quotient and a stable subobject of the same phase, each of class \mathbf{v}_0 .

¹Note that it is not an irreducible holomorphic symplectic variety when $\mathbf{v}^2 > 0$; instead it is deformation equivalent to $\text{Hilb}^{\mathbf{v}^2/2}(X) \times X$.

Proof. By the same argument as in the proof of [BMS16, Proposition A.8], the set of semistable objects of class \mathbf{v} is constant as σ varies. Hence we can assume σ is in the Gieseker-chamber for \mathbf{v}_0 and \mathbf{v} . The moduli space $M_\sigma(\mathbf{v}_0)$ is an abelian surface derived equivalent to X (up to a Brauer twist). The associated Mukai transform sends objects in $M_\sigma(\mathbf{v}_0)$ to skyscraper sheaves of points, and objects in $M_\sigma(\mathbf{v})$ to 0-dimensional torsion sheaves of length m . This proves the claim. \square

Finally, we need the construction of associated Kummer varieties. Let $\widehat{X} := \text{Pic}^0(X)$ be the dual abelian variety, and let \mathcal{P} be the Poincaré line bundle on $X \times \widehat{X}$. Let $\Phi^{\mathcal{P}}: D^b(X) \rightarrow D^b(\widehat{X})$ be the associated Fourier-Mukai transform. Given $E \in D^b(X)$, let $\det(E)$ be the associated determined line bundle. Now consider a moduli space $M_\sigma(\mathbf{v})$ with chosen basepoint E_0 . We obtain a map

$$(3) \quad \underline{\det}: M_\sigma(\mathbf{v}) \rightarrow \widehat{X}, \quad E \mapsto \det(E) \otimes \det(E_0)^\vee, \quad \text{and, dually,}$$

$$(4) \quad \widehat{\underline{\det}}: M_\sigma(\mathbf{v}) \rightarrow X, \quad E \mapsto \det(\Phi^{\mathcal{P}} E) \otimes \det(\Phi^{\mathcal{P}} E_0)^\vee.$$

Proposition 2.6 (Yoshioka). *Assume that $\mathbf{v}^2 \geq 2$ is primitive. Then the Albanese map of $M_\sigma(\mathbf{v})$ is given by*

$$\underline{\det} \times \widehat{\underline{\det}}: M_\sigma(\mathbf{v}) \rightarrow X \times \widehat{X},$$

and it has connected fibers.

If $\mathbf{v}^2 = 0$ and with \mathbf{v} primitive, then $M_\sigma(\mathbf{v})$ is an abelian surface. If moreover the rank $\text{rk}(\mathbf{v})$ is positive, then $\underline{\det}: M_\sigma(\mathbf{v}) \rightarrow \widehat{X}$ is a finite map of degree $\text{rk}(\mathbf{v})^2$.

Proof. For $\mathbf{v}^2 \geq 6$, this is part of the statement of [MY11, Proposition 5.16, (2a)]. For $\mathbf{v}^2 = 4$ and $\mathbf{v}^2 = 2$ this is proven for moduli spaces of Gieseker-stable sheaves in [Yos99a, Section 3] and [Yos99b, Corollary 4.3]. This property remains preserved under wall-crossing, and thus holds for any $M_\sigma(\mathbf{v})$, with the same proof as the one given in [MY11] for $\mathbf{v}^2 \geq 6$.

In case $\mathbf{v}^2 = 0$, then $M_\sigma(\mathbf{v})$ is (up to shift) a moduli space of semihomogeneous vector bundles. In particular, for $E \in M_\sigma(\mathbf{v})$ there is a map $\Pi: \widehat{X} \rightarrow M_\sigma(\mathbf{v})$, $L \mapsto E \otimes L$ that is surjective [Muk78, Proposition 6.10] of degree $\text{rk}(\mathbf{v})^2$ [Muk78, Proposition 7.1]. Since the composition $\underline{\det} \circ \Pi$ is multiplication by $\text{rk}(\mathbf{v})$, and thus of degree $\text{rk}(\mathbf{v})^4$, the claim follows. \square

If in fact $\mathbf{v}^2 \geq 4$, then the fibers of the Albanese map are irreducible holomorphic symplectic varieties (or K3 surfaces) of dimension $\mathbf{v}^2 - 2$, but we will not need that fact.

3. THE GIESEKER-WALL

The Gieseker chamber described by Proposition 2.4 typically has a wall corresponding to the Gieseker-Uhlenbeck contraction, corresponding to the subspace spanned by \mathbf{v} and $(0, 0, 1)$. In this section, we will describe its second wall for classes \mathbf{v} with $c_1(\mathbf{v}) = H$ and $\chi(\mathbf{v}) < 0$.

Proposition 3.1. *Let $\mathbf{v} = (r, H, \chi) \in H_{\text{alg}}^*(X, \mathbb{Z})$ with $r \geq 0$. For any $\alpha > 0$ and $\beta = 0$, we have $M_{\sigma_{\alpha,0}}(\mathbf{v}) = M_H(\mathbf{v})$. Similarly, all line bundles $L \in \text{Pic}^0(X)$ of degree 0 are $\sigma_{\alpha,0}$ -stable.*

Proof. Observe that

$$\begin{aligned}\Im Z_{\alpha,0}(E) &\in \mathbb{Z}_{\geq 0} \cdot \alpha H^2 \quad \text{for all } E \in \text{Coh}^\beta X, \text{ and} \\ \Im Z_{\alpha,0}(\mathbf{v}) &= \alpha H^2.\end{aligned}$$

Therefore, an object $E \in \text{Coh}^0 X$ with $\text{ch}(E) = \mathbf{v}$ can never be strictly $\sigma_{\alpha,0}$ -semistable: its Jordan-Hölder factors E_i would satisfy $\Im Z_{\alpha,0}(E_i) \in (0, \alpha H^2)$ in contradiction to the first equation. Combined with Propositions 2.2 and 2.4, this proves the first claim.

The stability of $L \in \text{Pic}^0(X)$ immediately follows from the last claim of Proposition 2.2, again combined with Proposition 2.4. \square




FIGURE 2. The space $M_\sigma(\mathbf{v})$ is unchanged in the chamber bounded by walls of $\overline{\mathbf{v}\mathcal{O}_p}$ and $\overline{\mathbf{v}\mathcal{O}_X}$.

The line connecting \mathbf{v} with $\text{Ker } Z_{0,0} = \mathbb{R} \cdot (1, 0, 0)$ is therefore the first possibility for a second wall of the Gieseker-chamber for \mathbf{v} . For $\chi < 0$, this wall does exist and will be described in the following.

Definition 3.2. (a) Given $\chi < 0$, let R be the number

$$R := \max \left\{ r \in \mathbb{Z}_{\geq 0} \mid (-r, H, \chi)^2 = H^2 + 2r\chi \geq 0 \right\} = \left\lfloor \frac{H^2}{-2\chi} \right\rfloor.$$

We denote w_0 by the character $(-R, H, \chi) \in H_{\text{alg}}^*(X, \mathbb{Z})$, and $w_k = w_0 + (k, 0, 0)$ for $k \in \mathbb{Z}_{\geq 0}$.

- (b) Let $\mathcal{W}_\chi \subset H_{\text{alg}}^*(X, \mathbb{R}) \cong \mathbb{R}^3$ be the two-dimensional subspace spanned by $(1, 0, 0)$ and w_0 (and containing all w_k). Denote by $\sigma_0 = (\text{Coh}^\beta X, Z_{\alpha, \beta})$ a stability condition on the wall corresponding to \mathcal{W}_χ : it is given as $\sigma_{\alpha_0, \beta_0}$ as in Theorem 2.1 for any α_0, β_0 with $\text{Ker } Z_{\alpha_0, \beta_0} \subset \mathcal{W}_\chi$.
- (c) Finally, let \mathcal{P}_0 denote the category of σ_0 -semistable objects in $\text{Coh}^{\beta_0} X$ of the same slope ν_0 as objects of character w_k (for any $k \geq 0$). The category \mathcal{P}_0 does not depend on the choice of σ_0 on \mathcal{W}_χ .

Given an object $E \in \mathcal{P}_0$, its Chern character is a linear combination

$$(5) \quad \chi(E) = a(1, 0, 0) + bw_0.$$

Since H generates the Picard group, b has to be integral, and therefore a as well. Moreover, $Z_{\alpha_0, \beta_0}(E)$ has to lie on the same ray as $Z_{\alpha_0, \beta_0}(w_k)$; combined with $\text{ch}(E)^2 \geq 0$ this is only possible if $b \geq 0$. This leads to the following observation:

Lemma 3.3. *Let $E \in \mathcal{P}_0$ be a σ_0 -semistable object with $\text{ch}(E) = w_k$. Then for each of its Jordan-Hölder factor E_i , the Chern character is either given by $\text{ch}(E_i) = w_{k'}$ for some $0 \leq k' \leq k$, or by $\text{ch}(E_i) = (1, 0, 0)$. Moreover, \mathcal{W}_χ is a wall for the Chern character w_k for all $k > 0$.*

Proof. Since $w_k - \text{ch}(E_i)$ is the sum of the characters of the remaining Jordan-Hölder factors, we must, in addition to the observations of the previous paragraph have $b \leq 1$.

If $b = 0$, i.e. $\text{ch}(E_i) = (a, 0, 0)$, then Proposition 2.5 shows $\text{ch}(E_i) = (1, 0, 0)$. If $b = 1$, we have $\text{ch}(E_i) = w_{k-a}$.

It remains to prove that \mathcal{W}_χ is a wall for w_k for $k > 0$. By Theorem 2.3 and the definition of w_k and R , there exists a stable object E' with Chern character w_{k-1} . Then $E' \oplus \mathcal{O}_X$ is a strictly σ_0 -semistable object, and so \mathcal{W}_χ describes a wall for the Chern character w_k . \square

Fix $K \geq 0$. Let σ_+ be a geometric stability condition, i.e. one given as in Theorem 2.1, sufficiently nearby the wall corresponding to \mathcal{W}_χ , on the side where the associated slope function satisfies $\nu_+(\mathcal{O}) < \nu_+(w_k)$ for $k \geq 0$. It is immediate from the local finiteness of walls (see Proposition 2.2) that σ_+ is not separated by a wall from \mathcal{W}_χ for any w_k and $0 \leq k \leq K$; in particular, $M_{\sigma_+}^s(w_k) = M_{\sigma_+}(w_k) \subset M_{\sigma_0}^{ss}(w_k)$. By Proposition 3.1, the stability condition σ_+ is in fact in the (dual of the) Gieseker-chamber of w_k . Moreover:

Proposition 3.4 ([BM14, Proposition 5.1]). *Let E be σ_0 -semistable and of Chern character w_k for $0 \leq k \leq K$. Let E_i be any of its Harder-Narasimhan filtration factors with respect to σ_+ . Then E_i is σ_0 -semistable, and its Chern character is contained in \mathcal{W}_χ . Moreover, the Jordan-Hölder factors of E_i are a subset of the Jordan-Hölder factors of E .*

Corollary 3.5. *Let $E \in \text{Coh}^{\beta_0}$ be a σ_0 -semistable object with character w_k . Then it is σ_+ -stable if and only if for every $L \in \text{Pic}^0(X)$ we have $\text{Hom}(E, L) = 0$.*

Proof. Since $\nu_+(L) = \nu_+(\mathcal{O}) < \nu_+(w_k)$, and since L is σ_+ -stable by Proposition 2.2, the condition is clearly necessary.

Conversely, assume that E is σ_0 -semistable but unstable with respect to σ . By Proposition 3.4, there is a destabilizing quotient $E \twoheadrightarrow E'$ such that E' is σ_0 -semistable with $\text{ch}(E') \in \mathcal{W}_\chi$, and such that the Jordan-Hölder factors of E' are a subset of those of E . By Lemma 3.3, the character of E' is either equal to $w_{k'}$ for $k' < k$, or equal to $(r, 0, 0)$. Since $\nu_+(E) \geq \nu_+(E')$, the former case is impossible, and so $\text{ch}(E') = (r, 0, 0)$. In light of Proposition 2.5, we can in fact assume $r = 1$, and so $E' = L$ for some $L \in \text{Pic}^0(X)$. \square

The following lemma is well-known.

Lemma 3.6. *Let E be an object that is σ_0 -semistable, and let S be one of its JH factors. Then there exists a unique short exact sequence*

$$T \hookrightarrow E \twoheadrightarrow E'$$

such that T is σ_0 -semistable with all JH factors isomorphic to S , and such that $\text{Hom}(S, E') = 0$.

Proof. The existence of E' is proved by induction on the length of the Jordan-Hölder filtration of E . If $\tilde{T} \hookrightarrow E \twoheadrightarrow \tilde{E}'$ is another such short exact sequence, then $\text{Hom}(T, \tilde{E}') = 0 = \text{Hom}(\tilde{T}, E')$, and a simple diagram-chase proves the uniqueness. \square

We will use this in the following context.

Definition 3.7. Let E be an object an object that is σ_0 -semistable. We apply Lemma 3.6 with $S = \mathcal{O}_X$ and define E_{red} as the unique object with a short exact sequence

$$(6) \quad T \hookrightarrow E \twoheadrightarrow E_{\text{red}},$$

such that T is an iterated self-extension of \mathcal{O}_X , and $\text{Hom}(\mathcal{O}_X, E_{\text{red}}) = 0$.

4. MAIN PROOF

We continue to use the notation from the previous section, in particular see Definition 3.2, and we continue to assume $\chi < 0$. The goal of this section is Theorem 4.12, on non-emptiness and dimension of the loci

$$M_k^h := \left\{ E \in M_{\sigma_+}^s(w_k) \mid \text{hom}(\mathcal{O}_X, E) = h \right\}.$$

The strategy is to control the existence of objects in M_k^h , and to construct them, via the two short exact sequences appearing in Lemma 4.2, respectively. First we observe the following:

Lemma 4.1. *If $E \in M_k^h$, then $\dim \text{Ext}^1(E, \mathcal{O}_X) = h - \chi$.*

Proof. This follows from Serre duality, $\chi(E) = \chi$, and $\text{Hom}(E, \mathcal{O}_X) = 0$ by stability. \square

Lemma 4.2. *Let E be an object in M_k^h .*

(a) Given a subspace $W \subset H^0(E)$, consider the short exact sequence in \mathcal{P}_0 given as

$$(7) \quad \mathcal{O}_X \otimes W \hookrightarrow E \rightarrow E'.$$

Then E' is σ_+ -stable with $h^0(E') \geq \dim W + \chi$.

(b) Conversely, let $V \subset \text{Ext}^1(E, \mathcal{O}_X)$, and consider the natural short exact sequence

$$(8) \quad \mathcal{O}_X \otimes V^\vee \hookrightarrow \tilde{E} \rightarrow E.$$

Then \tilde{E} is σ_+ -stable with $h^0(\tilde{E}) \geq \dim V$.

Proof. Since E' and \tilde{E} are σ_0 -semistable, in light of Corollary 3.5 we need to verify that they do not admit $L \in \text{Pic}^0(X)$ as a quotient in \mathcal{P}_0 . This is immediate for E' . For \tilde{E} , we consider the long exact sequence associated via $\text{Hom}(_, L)$:

$$0 = \text{Hom}(E, L) \rightarrow \text{Hom}(\tilde{E}, L) \hookrightarrow \text{Hom}(\mathcal{O}_X, L) \otimes V \rightarrow \text{Hom}(E, L[1]).$$

For $L \neq \mathcal{O}_X$, the vanishing is immediate. For $L = \mathcal{O}_X$, it follows by our choice of V .

The bound for $h^0(\tilde{E})$ is immediate, as W is a subspace of $H^0(\tilde{E})$ and the quotient injects into $H^0(E)$. As for E' , observe that $H^2(E') = \text{Hom}(E', \mathcal{O}_X)^\vee = 0$ and, due to the existence of the non-trivial extension E , also $\dim H^1(E') = \text{Ext}^1(E', \mathcal{O}_X) \geq \dim W$. Combined with $h^0(E') = \chi + h^1(E')$ this shows the remaining claim. \square

Their key difficulty is that for both short exact sequences above, we only obtain a bound for h^0 of the corresponding object, and that $h^0(E')$ may be non-zero even for $W = H^0(E)$. (In contrast, in the case of K3 surface, these dimensions are directly determined due to $h^1(\mathcal{O}_X) = 0$.) As we have explained previously, we need to proceed by induction. Moreover, in order to control the dimension of H^0 precisely for *some* of these extensions, it turns out that we have to prove more precise statements in our induction claim. These more precise claims are based on Lemma 3.6, by involving the class of E_{red} .

Corollary 4.3. *Let E be an object in M_k^h , then E_{red} given in Definition 3.7 is σ_+ -stable. In particular, $E_{\text{red}} \in M_{k_{\text{red}}}^0$ for some $k_{\text{red}} \in \mathbb{Z}_{\geq 0}$.*

Proof. This follows from the inductive construction of E_{red} in Lemma 3.6, and Lemma 4.2. \square

Definition 4.4. We define a locally closed subset $M_{k, k_{\text{red}}}^h$ of M_k^h by

$$M_{k, k_{\text{red}}}^h := \left\{ E \in M_k^h \mid E_{\text{red}} \in M_{k_{\text{red}}}^h \right\}.$$

Our induction process will be controlled by the following piece-wise linear function:

Definition 4.5. Let $\Delta_{KLM} : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$ be a function inductively defined as follows:

- $\Delta_{KLM}(t) := t$, when $0 \leq t \leq 1$;
- $\Delta_{KLM}(t) := \Delta_{KLM}(t-1) + t$ for $t \geq 1$.

It is a continuous piece-wise linear function with $\Delta_{KLM}(n) = \binom{n}{2}$ for $n \in \mathbb{Z}_{\geq 0}$; explicitly,

$$(9) \quad \Delta_{KLM}(t) = \left(t - \frac{\lfloor t \rfloor}{2}\right) (\lfloor t \rfloor + 1) \quad \text{for all } t \in \mathbb{R}_{\geq 0}.$$

Proposition 4.6. *The space $M_{k,k_{\text{red}}}^h$ is empty when*

$$(10) \quad \frac{k - k_{\text{red}}}{-\chi} < \Delta_{KLM}\left(\frac{h}{-\chi}\right).$$

Before the proof, let us use the statement of the Proposition 4.6 in order to illustrate the purpose of the function Δ_{KLM} . Consider the short exact sequence (7) for $W = H^0(E)$, and assume that $E \in M_{k,k_{\text{red}}}^h$ for a tuple k, k_{red}, h that satisfies (10). Then $E' \in M_{k',k_{\text{red}}}^{h'}$ for $k' = k - h$ and $h' \geq h + \chi$. Then the tuple k', k_{red}, h' also satisfies (10) precisely because of the functional equation satisfied by Δ_{KLM} .

Proof. When $k \leq k_{\text{red}} - \chi$, suppose there is an object $E \in M_{k,k_{\text{red}}}^h$ such that inequality (10) holds, i.e. $h \geq k - k_{\text{red}} + 1 \geq 1$. Consider the quotient E' defined by

$$(11) \quad \mathcal{O}_X^{\oplus h} \cong \mathcal{O}_X \otimes \text{Hom}(\mathcal{O}_X, E) \hookrightarrow E \twoheadrightarrow E'.$$

Note that, by Definition 3.7, $E_{\text{red}} \cong (E')_{\text{red}}$. But since $E' \in M_{k-h}$, we have $(E')_{\text{red}} \in M_{t'} = M_{k_{\text{red}}}$ for $t' \leq k - h \leq k_{\text{red}} - 1$, this leads the contradiction.

When $k > k_{\text{red}} - \chi$, we proceed by induction on k . Suppose there is an object $E \in M_{k,k_{\text{red}}}^h$ such that the (10) holds. In particular, we have $h > -\chi$. We again consider E' fitting into the short exact sequence (11). By Lemma 4.2, E' is σ_+ -stable with $h^0(E') \geq h + \chi$.

As before, $E_{\text{red}} \simeq (E')_{\text{red}}$, therefore, $M_{k-h,k_{\text{red}}}^{h'} \neq \emptyset$ for some $h' \geq h + \chi$. By induction on k ,

$$\frac{k - h - k_{\text{red}}}{-\chi} \geq \Delta_{KLM}\left(\frac{h'}{-\chi}\right) \geq \Delta_{KLM}\left(\frac{h + \chi}{-\chi}\right) = \Delta_{KLM}\left(\frac{h}{-\chi}\right) - \frac{h}{-\chi}.$$

This contradicts the assumption on k . □

Definition 4.7. We write $d(k, h)$ for the expected dimension of M_k^h , which is given as

$$d(k, h) := w_0^2 - 2k\chi + 2 + h\chi - h^2 = w_k^2 + 2 + h\chi - h^2.$$

Lemma 4.8. *Every irreducible component of M_k^h has dimension at least $d(k, h)$.*

Proof. This could be proved similarly to the case of line bundles on curves, see e.g. [ACGH85, Section IV.3]; however, to treat the cases $k \geq R$ (in which case $M_{\sigma_+}(w_k)$ parameterizes stable sheaves) and $k < R$ (in which case it instead parameterizes derived duals of stable sheaves) simultaneously, we present here a derived category version of the classical argument.

Consider any family $\mathcal{E} \in D^b(S \times X)$ of σ^+ -semistable objects of class w_k over a scheme S of finite type over \mathbb{C} ; this means that the derived restriction $\mathcal{E}_s := \mathcal{E}|_{\{s\} \times X} \in D^b(X)$ is an σ^+ -stable object of class w_k for every closed point s . We will prove that the locus

$$S_h := \{s \in S : h^0(\mathcal{E}_s) = h\} \subset S$$

has codimension at most $-h\chi + h^2$.

Let $\pi_S : S \times X \rightarrow S$ denote the projection, and consider the derived push-forward $\mathcal{F} := (\pi_S)_* \mathcal{E}$. By derived base change (see e.g. [Sta16, Tag 08IB]), we have $H^i(\mathcal{F} \otimes^{\mathbf{L}} \mathcal{O}_s) = H^i(\mathcal{E}_s) = 0$ for $i \neq 0, 1$ (note that $H^2(\mathcal{E}_s) = \text{Hom}(\mathcal{E}_s, \mathcal{O}_X)^\vee = 0$ since \mathcal{E}_s is σ_+ -stable). By [BM02, Proposition 5.4], it follows that \mathcal{F} is quasi-isomorphic to a 2-term complex of vector bundles $\mathcal{F}_0 \rightarrow \mathcal{F}_1$, and of rank $\chi = \text{rk } \mathcal{F}_0 - \text{rk } \mathcal{F}_1$. Then S_h is the locus where the rank of the differential is given by $\text{rk } \mathcal{F}_1 - h$; this has codimension at most $h(h - \chi)$ as claimed. \square

Remark 4.9. The proof evidently applies in much bigger generality: X could be an arbitrary scheme of finite type of the base field; we only need to assume that $\mathcal{E} \in D^b(S \times X)$ has the property that for all $s \in S$, the restriction \mathcal{E}_s is in $D^b(X)$, has compact support, and cohomology in at most two degrees. The classical proof instead constructs the two-term complex $\mathcal{F}_0 \rightarrow \mathcal{F}_1$ explicitly.

Let D be the remainder of division of h by $-\chi$, with $0 \leq D < -\chi$.

Proposition 4.10. *The moduli space $M_{k, k_{\text{red}}}^h$ is non-empty when $k \geq k_{\text{red}} \geq 0$ and h is the maximum integer such that*

$$(12) \quad \frac{k - k_{\text{red}}}{-\chi} \geq \Delta_{KLM} \left(\frac{h}{-\chi} \right).$$

When equality holds, $M_{k, k_{\text{red}}}^h$ is a bundle over $M_{k_{\text{red}}}^0$ of Grassmannians of D -dimensional subspaces in a $(-\chi)$ -dimensional vector space, and of total dimension $d(k, h)$.

Proof. When $k = k_{\text{red}}$, since $w_{k_{\text{red}}}^2 \geq 0$, the moduli space $M_{\sigma_+}(w_{k_{\text{red}}})$ is non-empty by Theorem 2.3. Since any $E \in M_{\sigma_+}(w_{k_{\text{red}}})$ has only finitely many Jordan-Hölder factors with respect to σ_0 , we can use the action of $\text{Pic}^0(X)$ to find an object that does not have \mathcal{O}_X as a factor, and in particular satisfies $\text{Hom}(\mathcal{O}_X, E) = 0$. Therefore, $M_{k_{\text{red}}, k_{\text{red}}}^0$ is non-empty. Moreover, it is an open dense subset of $M_{\sigma_+}^s(w_{k_{\text{red}}})$, and thus of expected dimension $d(k, 0)$.

Now consider the case $0 < k - k_{\text{red}} < -\chi$. Then $h = k - k_{\text{red}}$ satisfies equality in (12). Let E be an object in $M_{k_{\text{red}}, k_{\text{red}}}^0$. By Lemma 4.1, $\text{Ext}^1(E, \mathcal{O}_X)$ has dimension χ , and thus we can choose a subspace V of dimension h . We consider the short exact sequence

$$\mathcal{O}_X \otimes V^\vee \hookrightarrow \tilde{E} \twoheadrightarrow E$$

as in Lemma 4.2. By the Lemma, \tilde{E} is σ_+ -stable. Clearly $(\tilde{E})_{\text{red}} = E$ and $H^0(\tilde{E}) = V^\vee$. Therefore, $\tilde{E} \in M_{k, k_{\text{red}}}^h$, proving the non-emptiness as claimed.

Our construction depended on a choice of a point in the Grassmannian bundle over $M_{k_{\text{red}}, k_{\text{red}}}^0$ whose fiber over E is given by the set of h -dimensional (or, equivalently, D -dimensional) subspaces in $\text{Ext}^1(E, \mathcal{O}_X) \cong \mathbb{C}^{-\chi}$. On the other hand, applying Lemma 4.2, part a, we see that any object $F \in M_{k, k_{\text{red}}}^h$ fits into a short exact sequence

$$\mathcal{O}_X \otimes H^0(F) \hookrightarrow F \twoheadrightarrow F_{\text{red}}$$

with $F_{\text{red}} \in M_{k_{\text{red}}, k_{\text{red}}}^0$, and thus this Grassmannian bundle describes the entire stratum $M_{k, k_{\text{red}}}^h$. Its dimension is

$$\begin{aligned} \dim M_{k_{\text{red}}}^0 + \dim Gr(k - k_{\text{red}}, -\chi) &= w_0^2 + 2 - 2k_{\text{red}}\chi + (k - k_{\text{red}})(-\chi - k + k_{\text{red}}) \\ &= w_0^2 + 2 - 2k\chi + (k - k_{\text{red}})(\chi - k + k_{\text{red}}) \\ &= w_0^2 + 2 - 2k\chi - (h - \chi)h \\ &= d(k, h). \end{aligned}$$

For $k - k_{\text{red}} \geq -\chi$, we proceed by induction on k . Let h be the integer as in the statement of the Proposition. Let h' be the maximum integer, such that

$$\Delta_{KLM} \left(\frac{h'}{-\chi} \right) \leq \frac{k - h - k_{\text{red}}}{-\chi}.$$

By the functional equation of Δ_{KLM} , $h' \geq h + \chi \geq 0$. By induction on k , there is an object E in $M_{k-h, k_{\text{red}}}^{h'}$. By Lemma 4.1 the dimension of $\text{Ext}^1(E, \mathcal{O}_X)$ is $h' - \chi \geq h$, and thus we can choose a subspace V of dimension h . We again consider the extension

$$\mathcal{O}_X \otimes V^\vee \hookrightarrow \tilde{E} \twoheadrightarrow E;$$

applying Lemma 4.2 as before shows that \tilde{E} is σ_+ -stable with $\left(\tilde{E} \right)_{\text{red}} \cong E_{\text{red}}$. In particular, $\tilde{E} \in M_{k, k_{\text{red}}}^{h''}$ for some $h'' \geq h$. Now since h is the maximum number satisfying (12), Proposition 4.6 says that $M_{k, k_{\text{red}}}^{h''}$ is empty for $h'' > h$. Hence $h'' = h$, and so $M_{k, k_{\text{red}}}^h$ is non-empty as claimed.

When equality holds, it remains to show the statement about the dimension. In that case, we have $h' = h + \chi$ in each induction step, and in particular the remainder D remains preserved at each step. We have a bijective morphism

$$\begin{aligned} M_{k, k_{\text{red}}}^h &\rightarrow M_{k-h, k_{\text{red}}}^{h+\chi} \\ E &\mapsto E' = E / (\mathcal{O}_X \otimes \text{Hom}(\mathcal{O}, E)); \end{aligned}$$

indeed, its inverse is given by associating to E the extension \tilde{E} given by

$$\mathcal{O}_X \otimes \text{Hom}(E, \mathcal{O}_X[1])^\vee \hookrightarrow \tilde{E} \twoheadrightarrow E.$$

It follows by induction that $M_{k, k_{\text{red}}}^h$ is a Grassmannian-bundle over $M_{k_{\text{red}}}^0$ of dimension

$$\begin{aligned} d(k - h, h + \chi) &= w_0^2 - 2(k - h)\chi + 2 + (h + \chi)\chi - (h - \chi)^2 \\ &= w_0^2 - 2k\chi + 2 + h\chi - h^2 = d(k, h). \end{aligned}$$

□

Lemma 4.11. *Let $h, k, k_{\text{red}} \in \mathbb{Z}_{\geq 0}$. Whenever $M_{k, k_{\text{red}}}^h$ is non-empty, it satisfies*

$$\dim M_{k, k_{\text{red}}}^h \leq d(k, h).$$

Moreover, if equality holds, then h, k, k_{red} satisfy equality in equation (12).

Proof. We prove the statement by induction on k , the case $k = 0$ being obvious. Similarly, for $h = 0$, we have $M_{k, k}^0 = M_k^0$, which is an open subset of $M_{\sigma_+}(w_k)$ by Proposition 4.10.

We thus assume $k > 0$ and $h > 0$. Given $E \in M_{k, k_{\text{red}}}^h$, again consider the quotient E' (in \mathcal{P}_0) defined by

$$\mathcal{O}_X \otimes H^0(E) \hookrightarrow E \twoheadrightarrow E',$$

which is σ_+ -stable with $h^0(E') \geq h + \chi$ by Lemma 4.2. We consider the map

$$\pi: M_{k, k_{\text{red}}}^h \rightarrow \bigcup_{l \geq \max(0, h + \chi)} M_{k-h, k_{\text{red}}}^l, \quad E \mapsto E',$$

where the right-hand-side is considered as a locally closed subset of $M_{\sigma_+}(w_{k-h})$.

It is sufficient to prove the dimension bound for each of the pre-images of the finitely many strata on the right-hand-side. By the induction on k , the dimension of $M_{k-h, k_{\text{red}}}^l$ does not exceed $d(k-h, l)$. For each $E' \in M_{k-h, k_{\text{red}}}^l$, we have $\dim \text{Ext}^1(E', \mathcal{O}_X) = l - \chi$ by Lemma 4.1. Therefore, $\pi^{-1}(E')$ is a locally closed subset of the Grassmannian of h -dimensional subspaces in \mathbb{C}^l (defined by the associated extension having h -dimensional space of global sections); in particular its dimension is at most $h(l - \chi - h)$.

Thus, if the pre-image of $M_{k-h, k_{\text{red}}}^l$ is non-empty, its dimension is bound by

$$\begin{aligned} \dim \pi^{-1} \left(M_{k-h, k_{\text{red}}}^l \right) &\leq \dim M_{k-h, k_{\text{red}}}^l + h(l - \chi - h) \\ &\leq d(k-h, l) + h(l - \chi - h) \\ &= d(k, h) + 2h\chi - l(l - \chi) + h(h - \chi) + h(l - \chi - h) \\ &= d(k, h) - l(l - \chi - h) \\ (13) \quad &\leq d(k, h). \end{aligned}$$

The last inequality achieves equality only if $l = 0 \geq h + \chi$ or $l = h + \chi > 0$.

In the former case, $M_{k-h, k_{\text{red}}}^0$ is non-empty only if $k_{\text{red}} = k - h$. Since $h + \chi \leq 0$, we have $\Delta_{KLM} \left(\frac{h}{-\chi} \right) = \frac{h}{-\chi} = \frac{k - k_{\text{red}}}{-\chi}$, and thus equality in (12).

In the second case $l = h + \chi$, by the induction on k , the second inequality above can only be an equality if $h + \chi, k - h, k_{\text{red}}$ satisfy equality in (12). By the functional equation of Δ_{KLM} , we conclude

$$\Delta_{KLM} \left(\frac{h}{-\chi} \right) = \Delta_{KLM} \left(\frac{h + \chi}{-\chi} \right) + \frac{h}{-\chi} = \frac{k - h - k_{\text{red}}}{-\chi} + \frac{h}{-\chi} = \frac{k - k_{\text{red}}}{-\chi}$$

as claimed. \square

Theorem 4.12. *The moduli space M_k^h is nonempty if and only if*

$$\frac{k}{-\chi} \geq \Delta_{KLM} \left(\frac{h}{-\chi} \right).$$

Whenever non-empty, M_k^h is irreducible with expected dimension $d(k, h)$.

Proof. Observe either by induction, or by the explicit formula (9), that $\chi \cdot \Delta \left(\frac{h}{-\chi} \right)$ is an integer. Therefore, given h and k satisfying the assumptions of the Theorem, there is a unique integer k_0 with $0 \leq k_0 \leq k$ such that equality holds in equation 12. By Proposition 4.10, for this choice of k_{red} , the stratum $M_{k, k_{\text{red}}}^h$ is non-empty, irreducible and of expected dimension $d(k, h)$.

On the other hand, for all other choices of k_{red} , Lemma 4.11 shows that the stratum $M_{k, k_{\text{red}}}^h$ has dimension strictly smaller than $d(k, h)$. By Lemma 4.8, it is thus contained in the closure of a stratum of bigger dimensions.

Conversely, if $\frac{k}{-\chi} < \Delta_{KLM} \left(\frac{h}{-\chi} \right)$, then Proposition 4.6 shows that M_k^h is empty. \square

Note that all constructions in this sections have been invariant under the action of X by translation. We note the following consequence of this fact when combined with the proof of Theorem 4.12:

Proposition 4.13. *Assume that M_k^h is non-empty, and let k_0 with $0 \leq k_0 \leq k$ be such that*

$$\frac{k - k_0}{-\chi} = \Delta_{KLM} \left(\frac{h}{-\chi} \right).$$

Let $0 \leq D < -\chi$ be the remainder of division of h by $-\chi$. Then M_k^h has a dense open subset M_{k, k_0}^h , invariant under the action of X by translation, that is isomorphic to a bundle over $M_{k_0}^0$ of Grassmannians of type $\text{Gr}(D, -\chi)$. Its fiber over a point $F \in M_{k_0}^0$ parameterises iterated extensions of F by \mathcal{O}_X .

5. CONCLUSION

We will now prove our main result, which we first recall:

Theorem 1.1. *Assume $\chi \neq 0$. The Brill-Noether locus $V_d^r(|H|)$ is non-empty if and only if*

$$(14) \quad \rho + g - 2 \geq D |\chi| - D^2,$$

where D denotes the remainder of division of $r + 1$ by $|\chi|$. Moreover, when it is non-empty, it is generically smooth and of expected dimension $\rho + g - 2$.

When the above inequality is strict, then $V_d^r(|H|)$ is irreducible. Otherwise, it is a disjoint union of Grassmannians.

Proof. Using the derived dual on X , or, equivalently, Serre duality on the individual curves, we first reduce to the case $\chi < 0$.

Thus we are in the situation of the previous section. By Lemma 3.1, we have $M_H(0, H, \chi) = M_{\sigma_+}^s(0, H, \chi) = M_{\sigma_+}(w_R)$. (Note that by Definition 3.2, the value R is non-negative by $\chi < 0$.)

Hence $V_d^r(|H|)$ is the intersection of the Brill-Noether variety M_R^{r+1} studied in the previous section with the set of sheaves supported on a curve in the linear system $|H|$ (rather than a translate of such a curve). If we consider the action of X on $M_H(0, H, \chi) = M_{\sigma_+}(w_R)$ by translation, then each orbit contains finitely many sheaves supported on a curve in the linear system $|H|$. Moreover, the action leaves the Brill-Noether variety M_R^{r+1} invariant.

It follows that $V_d^r(|H|)$ is non-empty if and only if M_R^{r+1} is non-empty, with

$$\begin{aligned} \dim V_d^r(|H|) &= \dim M_R^{r+1} - 2 = d(R, r+1) - 2 \\ &= (0, H, \chi)^2 + 2 + (r+1)\chi - (r+1)^2 - 2 \\ &= 2g - 2 - (r+1)(r+1-\chi) = \rho + g - 2. \end{aligned}$$

Let $r+1 = s(-\chi) + D$, where $s \in \mathbb{Z}_{\geq 0}$, $0 \leq D < \chi$. By Theorem 4.12, M_R^{r+1} is non-empty if and only if

$$\begin{aligned} \frac{R}{-\chi} &\geq \left(\frac{r+1}{-\chi} - \frac{1}{2} \left\lfloor \frac{r+1}{-\chi} \right\rfloor \right) \left(\left\lfloor \frac{r+1}{-\chi} \right\rfloor + 1 \right) \\ &\iff R \geq \frac{1}{2}(r+1+D)(s+1). \end{aligned}$$

The bound on the right is an integer: indeed, when s is even, $r+1$ and D have the same parity. Thus, we may replace R by $\frac{H^2}{-2\chi}$, omitting the round-down (see Definition 3.2). Therefore, M_R^{r+1} and $V_d^r(|H|)$ are non-empty if and only if

$$\begin{aligned} 2g - 2 - (r+1+D)(r+1-D-\chi) &\geq 0 \\ \iff 2g - 2 - (r+1)(r+1-\chi) + D^2 + D\chi &\geq 0 \\ \iff \rho + g - 2 &\geq D(-\chi) - D^2. \end{aligned}$$

Now assume that the inequality (14) is strict; we need to show that $V_d^r(|H|)$ is irreducible. Note that we can define $V_d^r(|H|)$ equivalently as the preimage of $0 \in \widehat{X}$ under the map

$$(15) \quad D: M_R^{r+1} \rightarrow \widehat{X}, \quad E \mapsto \det(E) \otimes \mathcal{O}(-H).$$

Now let k_0 be as in Proposition 4.13, and consider the open subset $M_{R,k_0}^{r+1} \subset M_R^{r+1}$. Since this inclusion is translation-invariant, it suffices to show that the restriction factors as

$$M_{R,k_0}^{r+1} \rightarrow M_{k_0}^0 \xrightarrow{\det \times \widehat{\det}} \widehat{X} \times X \rightarrow \widehat{X};$$

here the first map is the projection to the base of the Grassmannian bundle given in the Proposition 4.13, the second map is the restriction of the Albanese map appearing in Proposition 2.6, and the third one is, up to shift, given by projection to the first factor.

We claim that under our assumptions, each of these maps has connected fibers. Indeed, this is obvious for the first and the third map. For the second map, note that by Proposition 4.13, the dimension of M_{k_0} must be equal to

$$\dim M_{k_0} = \dim M_R^{r+1} - \dim \operatorname{Gr}(D, -\chi) = \rho + g - D(-\chi - D).$$

Thus, if the inequality (14) is strict, then M_{k_0} has dimension at least 4, and we can apply Proposition 2.6 to deduce connectedness of fibers for $\underline{\det} \times \widehat{\underline{\det}}$.

The composition is thus a map between smooth irreducible varieties with connected fibers; since the generic fiber is smooth and connected, it is irreducible. Since translation acts transitively on the set of fibers, it follows that M_{R,k_0}^{r+1} is smooth and irreducible.

Otherwise, if equality holds in (14), then M_{k_0} is two-dimensional, i.e. $w_{k_0}^2 = 0$. Hence we have $-k_0 = \operatorname{rk}(w_{k_0}) = \frac{H^2}{2\chi} = \frac{g-1}{\chi}$. By Proposition 2.6, the moduli space M_{k_0} contains k_0^2 many elements with determinant $\mathcal{O}(H)$ (rather than one of its translates). The variety $V_d^r(|H|)$ consists of one Grassmannian for each of these objects. \square

Remark 5.1. Since $0 \leq D < -\chi$, the right-hand-side is always positive, as we should expect. It's a little more tricky to see that if the right-hand side is bigger than $g - 2$ and if ρ violates the inequality, then in fact $\rho < 0$.

Remark 5.2. This inequality gives exactly the same bound as that in Theorem A.1 in [KLCM15]:

$$(16) \quad \rho + r(r+2) \geq -\left\lfloor \frac{r}{-\chi} \right\rfloor \chi \left(r+1 + \frac{1}{2} \chi \left(\left\lfloor \frac{r}{-\chi} \right\rfloor + 1 \right) \right).$$

When $\left\lfloor \frac{r}{-\chi} \right\rfloor = \left\lfloor \frac{r+1}{-\chi} \right\rfloor = s$, the inequality (16) is equivalent to

$$\begin{aligned} \rho + r(r+2) &\geq -s\chi(r+1 - \frac{1}{2}(-\chi)(s+1)) \\ \iff 2\rho + 2r(r+2) &\geq 2(r+1-D)(r+1 - \frac{1}{2}(r+1-D) - \frac{1}{2}\chi) \\ \iff \rho + g + (r+1)(\chi + r+1) - 2 &\geq (r+1-D)(r+1+D+\chi) \\ \iff \rho + g - 2 &\geq D(-\chi) - D^2. \end{aligned}$$

In the remaining case $r+1 = s(-\chi)$, the inequality (16) is equivalent to

$$\begin{aligned} \rho + r(r+2) &\geq (r+1+\chi)(r+1 + \frac{1}{2}\chi s) \\ \iff 2\rho + 2r(r+2) &\geq (r+1+\chi)(r+1) \\ \iff \rho + g - 2 &\geq 0. \end{aligned}$$

\square

Proof of Theorem 1.2. When $k < -R$, we have $\mathbf{v}^2 < 0$. The inequality fails since $\mathbf{v}^2 - (r+1)(r+1-\chi) < 0 \leq D(-\chi) - D^2$. The moduli space $M_H^{r+1}(v)$ is empty due to Proposition 2.5.

When $k \geq -R$, by Lemma 3.1, we have $M_H(k, H, \chi) = M_{\sigma_+}^s(k, H, \chi) = M_{\sigma_+}(w_{k+R})$. It follows that $M_H^{r+1}(v) = M_{k+R}^{r+1}$. We apply Theorem 4.12 for the space M_{k+R}^{r+1} . This space is non-empty if and only if

$$\begin{aligned} \frac{k+R}{-\chi} &\geq \left(\frac{r+1}{-\chi} - \frac{1}{2} \left\lfloor \frac{r+1}{-\chi} \right\rfloor \right) \left(\left\lfloor \frac{r+1}{-\chi} \right\rfloor + 1 \right) \\ \iff R &\geq \frac{1}{2}(r+1+D) \left(\frac{r+1-D}{-\chi} + 1 \right) - k. \end{aligned}$$

Note that the bound on the right is an integer, we may substitute this bound to the constraint on R . The space M_{k+R}^{r+1} is non-empty if and only if

$$\begin{aligned} 2g - 2 - (r+1+D) \left(\frac{r+1-D}{-\chi} + 1 \right) (-\chi) + 2k\chi &\geq 0 \\ \iff \mathbf{v}^2 + (r+1)\chi - (r+1)^2 &\geq D(-\chi) - D^2. \end{aligned}$$

By Theorem 4.12, when M_{R+k}^{r+1} is non-empty, it is irreducible of the expected dimension:

$$d(k+R, r+1) = \mathbf{v}^2 + 2 - (r+1)(r+1-\chi).$$

□

6. GENERALISATION

Let (X, H) be a polarized abelian surface satisfying Assumption (*). In this section, we explain how to adapt all our arguments from abelian Picard rank one to X .

Let $\Lambda_H \cong \mathbb{Z}^3$ denote the image of the map

$$v_H: K(X) \rightarrow \mathbb{R}^3, \quad E \mapsto (\text{ch}_0(E), H \cdot \text{ch}_1(E), \text{ch}_2(E)).$$

We will only consider stability conditions for which the central charge factors via v_H , and denote the space of such stability conditions by $\text{Stab}_H(X)$. The pair $\sigma_{\alpha, \beta} := (\text{Coh}^\beta X, Z_{\alpha, \beta})$ defines a stability condition on $D^b(X)$ and there is a continuous map from $\mathbb{R}_{>0} \times \mathbb{R} \rightarrow \text{Stab}(X)$. The only difference here is that $\text{Stab}_H(X)$ is not the whole space but a slice of the whole space. The slope function $\nu_{\alpha, \beta}$ is defined in the same way.

The Mukai pairing equips $\Lambda_H \otimes \mathbb{R}$ with the quadratic form Q , and identifies the upper half plane $\mathbb{R}_{>0} \times \mathbb{R}$ with the projectivization of the negative cone of Q , via taking the kernel of $Z_{\alpha, \beta}$ in $\Lambda_H \otimes \mathbb{R}$. All the propositions in Section 2 hold for the higher Picard rank case.

In the proof of Proposition 3.1, we use the fact that ‘ $\Im Z_{\alpha, 0}(E)$ is of the form $\mathbb{Z}_{\geq 0} \cdot H^2$ for all $E \in \text{Coh}^\beta X$ ’. This also holds by Assumption (*). For Lemma 3.3, we shall modify the statement to ‘ $v_H(E_i)$ is either $v_H(w_{k'})$ or $(r, 0, 0)$ ’, and prove it in the same way as the Picard number one case. Similarly, for Corollary 3.5, we change the condition to ‘for any σ_0 -stable object O such that

$v_H(O) = (r, 0, 0)'$. Note that this modification does not affect the argument for Lemma 4.2, which is the only place that uses Corollary 3.5. All the other statements do not rely on the Picard rank.

REFERENCES

- [ACGH85] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris. *Geometry of algebraic curves. Vol. I*, volume 267 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, 1985.
- [Bay16] Arend Bayer. Wall-crossing implies Brill-Noether. Applications of stability conditions on surfaces, 2016, arXiv:1604.08261.
- [BM02] Tom Bridgeland and Antony Maciocia. Fourier-Mukai transforms for $K3$ and elliptic fibrations. *J. Algebraic Geom.*, 11(4):629–657, 2002, arXiv:math/9908022.
- [BM14] Arend Bayer and Emanuele Macrì. MMP for moduli of sheaves on $K3$ s via wall-crossing: nef and movable cones, Lagrangian fibrations. *Invent. Math.*, 198(3):505–590, 2014, arXiv:1301.6968.
- [BMS16] Arend Bayer, Emanuele Macrì, and Paolo Stellari. The space of stability conditions on abelian threefolds, and on some Calabi-Yau threefolds, 2016, arXiv:1410.1585.
- [Bri08] Tom Bridgeland. Stability conditions on $K3$ surfaces. *Duke Math. J.*, 141(2):241–291, 2008, arXiv:math/0307164.
- [HRS96] Dieter Happel, Idun Reiten, and Sverre O. Smalø. Tilting in abelian categories and quasitilted algebras. *Mem. Amer. Math. Soc.*, 120(575):viii+ 88, 1996.
- [JR17] David Jensen and Dhruv Ranganathan. Brill-Noether theory for curves of a fixed gonality, 2017, arXiv:1701.06579.
- [Kem71] G. Kempf. *Schubert Methods with an Application to Algebraic Curves [by] G. Kempf*. Mathematisch Centrum, Amsterdam. Afdeling Zuivere Wiskunde [Publications] 71-6. 1971.
- [KL72] Steven L. Kleiman and Dan Laksov. On the existence of special divisors. *Amer. J. Math.*, 94:431–436, 1972.
- [KLCM15] Andreas Leopold Knutsen, Margherita Lelli-Chiesa, and Giovanni Mongardi. Severi varieties and Brill-Noether theory of curves on abelian surfaces, 2015, arXiv:1503.04465.
- [Laz86] Robert Lazarsfeld. Brill-Noether-Petri without degenerations. *J. Differential Geom.*, 23(3):299–307, 1986.
- [Mac12] Antony Maciocia. Computing the walls associated to Bridgeland stability conditions on projective surfaces. *Asian J. Math.*, 18(2):263–279, 2012, arXiv:1202.4587. To appear in *Asian J. Math.*
- [MS16] Emanuele Macrì and Benjamin Schmidt. Introduction to Bridgeland stability, 2016. Lecture notes for the CIMP-ICTP-Mexico research school 2016. In preparation.
- [Muk78] Shigeru Mukai. Semi-homogeneous vector bundles on an Abelian variety. *J. Math. Kyoto Univ.*, 18(2):239–272, 1978.
- [MY11] Hiroki Minamide, Shintarou Yanagida, and Kōta Yoshioka. Some moduli spaces of Bridgeland’s stability conditions. *Int. Math. Res. Not. IMRN*, (19):5264–5327, 2011.
- [Pf13] Nathan Pflueger. On linear series with negative Brill-Boether number, 2013, arXiv:1311.5845.
- [Pf17] Nathan Pflueger. Brill–Noether varieties of k -gonal curves. *Advances in Mathematics*, 312:46–63, 2017.
- [Sta16] The Stacks Project Authors. Stacks project, 2016. Available at <http://stacks.math.columbia.edu>.
- [Yos99a] Kota Yoshioka. Albanese map of moduli of stable sheaves on abelian surfaces, 1999, arXiv:math.AG/9901013.
- [Yos99b] Kōta Yoshioka. Some notes on the moduli of stable sheaves on elliptic surfaces. *Nagoya Math. J.*, 154:73–102, 1999.
- [Yos12] Kōta Yoshioka. Bridgeland’s stability and the positive cone of the moduli spaces of stable objects on an abelian surface, 2012, arXiv:1206.4838.

SCHOOL OF MATHEMATICS AND MAXWELL INSTITUTE, UNIVERSITY OF EDINBURGH, JAMES CLERK MAXWELL BUILDING, PETER GUTHRIE TAIT ROAD, EDINBURGH, EH9 3FD, UNITED KINGDOM

E-mail address: arend.bayer@ed.ac.uk

URL: <http://www.maths.ed.ac.uk/~abayer/>

SCHOOL OF MATHEMATICS AND MAXWELL INSTITUTE, UNIVERSITY OF EDINBURGH, JAMES CLERK MAXWELL BUILDING, PETER GUTHRIE TAIT ROAD, EDINBURGH, EH9 3FD, UNITED KINGDOM

E-mail address: chunyi.li@ed.ac.uk

URL: <https://sites.google.com/site/chunyi0401/>